

Unless otherwise stated, use a coordinate system with $+x$ to the right and $+y$ upward, toward the top edge of the page.

Section 1.

Questions 1–2: Copper has a density of 8960 kg/m^3 and an atomic mass of 63.55 g/mol . One atom of copper has 29 protons and 29 electrons. Avogadro’s number is $6.02 \times 10^{23} \text{ atoms/mol}$. A certain copper BB has a volume $5.00 \times 10^{-10} \text{ m}^3$.

1. How many electrons are in the BB?

- (a) 1.23×10^{21} electrons
- (b) 4.24×10^{19} electrons
- (c) 2.46×10^{19} electrons
- (d) 2.70×10^{18} electrons
- (e) 7.82×10^{19} electrons

2. If the BB has an electric charge of -2 nC , how many additional electrons did it gain?

- (a) 3.2×10^{28} electrons
- (b) 1.25×10^{10} electrons
- (c) 1.25×10^{19} electrons
- (d) 2.0×10^9 electrons
- (e) 8.0×10^{11} electrons

3. Two equal magnitude charges are shown in four different configurations.




© 2010 Pearson Education, Inc.

In which case(s) does the *net* electric field at the dot point to the left.

- (a) A only
- (b) D only
- (c) C only
- (d) C and D
- (e) A, C, and D


Questions 4-5: A proton $+q$ is near a neutral atom (the circle) as shown below.



4. What is the direction of the force by the proton on the neutral atom?

- (a) to the right, $+x$
- (b) to the left, $-x$
- (c) upward, $+y$
- (d) downward, $-y$
- (e) none of the above because it is zero

5. Which of these diagrams shows the polarization of the neutral atom?



- (a) A
- (b) B
- (c) C
- (d) D

Questions 6-8: You have four metal disks with charges: $q_A = -8.0 \mu\text{C}$, $q_B = -2.0 \mu\text{C}$, $q_C = +5.0 \mu\text{C}$, and $q_D = +12.0 \mu\text{C}$.

6. Holding them with insulators, you bring two disks together so that they touch. Then you separate them. You measure the resulting charge of each disk and find that it is $+5.0 \mu\text{C}$ per disk. Which two disks did you bring together?

- (a) A and B
- (b) B and C
- (c) C and D
- (d) B and D
- (e) A and D


7. If you touch disk A and disk B together,

- (a) A will gain electrons
- (b) A will gain protons
- (c) A will lose electrons
- (d) A will lose protons

8. If you touch disk C and disk D together,

- (a) C will gain electrons
- (b) C will gain protons
- (c) C will lose electrons
- (d) C will lose protons

Questions 9–10: Use this image to answer each of the following questions.




9. Which sphere is a neutral insulator in an applied electric field (due to other charges not shown) that is to the right?

- (a) A
- (b) B
- (c) C
- (d) D
- (e) O

10. Which sphere is a neutral conductor in an applied electric field (due to other charges not shown) that is to the left?


- (a) M
- (b) N
- (c) C
- (d) D
- (e) O

Questions 11–13: Three charged spheres are arranged as shown below. In the figure, $d = 4 \text{ cm}$ and $q = 1 \mu\text{C}$.



11. What is the magnitude of the force by the particle of charge $-2q$ on the particle of charge $-q$?
 - (a) 5.6 N
 - (b) 0.45 N
 - (c) 2.8 N
 - (d) 22 N
 - (e) 11 N
12. What is the magnitude of the *net* force on the particle of charge $-q$?
 - (a) zero
 - (b) 17 N
 - (c) 11 N
 - (d) 22 N
 - (e) 5.6 N
13. What is the direction of the *net* force on the particle of charge $-q$?
 - (a) to the right, $+x$
 - (b) to the left, $-x$
 - (c) none of the above because it is zero


Questions 14–16: A dipole made of charged spheres with $q = 8 \text{ nC}$, $d = 4 \text{ cm}$ is shown below?



14. What is the magnitude of the electric field due to $-q$ at the location of the dot?
 - (a) 45,000 N/C
 - (b) 80,000 N/C
 - (c) 57,600 N/C
 - (d) 90,000 N/C
 - (e) 28,800 N/C
15. What is the angle with respect to the horizontal for the electric field due to $-q$ at the location of the dot?
 - (a) 37°
 - (b) 45°
 - (c) 30°
 - (d) 53°
 - (e) 60°

16. What is the magnitude of the *net* electric field at the location of the dot?
- (a) 97,300 N/C
 - (b) 109,000 N/C
 - (c) 120,000 N/C
 - (d) 100,000 N/C
 - (e) 66,800 N/C

Questions 17–19: In an experiment, an electron travels near a nucleus along the path shown below.




17. The potential difference, $\Delta V = V_f - V_i$, is
- (a) positive.
 - (b) negative.
 - (c) zero.
18. As the electron travels from *i* to *f*, the potential energy, U , of the system (of electron and nucleus)
- (a) increases.
 - (b) decreases.
 - (c) remains constant.
19. As it travels from *i* to *f*, the kinetic energy of the electron
- (a) increases.
 - (b) decreases.
 - (c) remains constant.

Questions 20–21: In a particular LEGO kit, a capacitor with a capacitance of 0.5 F is charged by a solar cell. The solar cell produces a potential difference of 4 V across the capacitor. The area of each plate is 0.01 m² and they are separated 0.1 mm.

20. How much charge is stored on each capacitor plate?
- (a) 0.125 C
 - (b) 0.5 C
 - (c) 2.0 C
 - (d) 8.0 C
 - (e) 4.0 C
21. What is the dielectric constant K of the insulator that is between the plates?
- (a) 1.1×10^{13}
 - (b) 2.3×10^9
 - (c) 4.5×10^9
 - (d) 1.1×10^9
 - (e) 5.6×10^8

Questions 22–25: Two electrodes, shown below, are connected to a battery with a potential difference of 1 V. The dotted lines are 1 cm apart.



22. What is the magnitude of the electric field between the plates?
- (a) 4 V/m
 - (b) 50 V/m
 - (c) 25 V/m
 - (d) 100 V/m
23. What is the direction of the electric field between the plates?
- (a) to the right
 - (b) to the left
 - (c) upward
 - (d) downward
 - (e) none of the above because it is zero
24. What is the electric potential V at point B?
- (a) 0 V
 - (b) 0.25 V
 - (c) 0.5 V
 - (d) 0.75 V
 - (e) 1.0 V
25. If an electron travels from point A to point B, what is its change in electric potential energy, ΔU , in eV?
- (a) 0
 - (b) 0.25 eV
 - (c) 0.5 eV
 - (d) 0.75 eV
 - (e) 1.0 eV