

# A modified version of “Preparation of manuscripts for the American Journal of Physics using L<sup>A</sup>T<sub>E</sub>X ” by Harvey Gould

modified by Aaron Titus

November 30, 2010

## 1 Introduction

T<sub>E</sub>X looks more difficult than it is. It is almost as easy as  $\pi$ . See how easy it is to make special symbols such as  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\sin x$ ,  $\hbar$ ,  $\lambda$ ,  $\dots$ . We also can make subscripts  $A_x$ ,  $A_{xy}$  and superscripts,  $e^x$ ,  $e^{x^2}$ , and  $e^{ab}$ . We will use L<sup>A</sup>T<sub>E</sub>X, which is based on T<sub>E</sub>X and has many higher-level commands (macros) for formatting, making tables, etc. More information can be found in Ref. [1], a book written by Kopka and Daly.[1]

We just made a new paragraph. Extra lines and spaces make no difference. Note that all formulae are enclosed by  $\$$  and occur in *math mode*.

The default font is Computer Modern. It includes *italics* or *italics*, **boldface** or **boldface**, *slanted* or *slanted*, and **monospaced** or **monospaced** (typewriter) fonts.

## 2 Equations

Let us see how easy it is to write equations.

$$\Delta = \sum_{i=1}^N w_i (x_i - \bar{x})^2. \quad (1)$$

For AJP all equations must be numbered, however for other uses, we can have an equation without a number by writing

$$P(x) = \frac{x - a}{b - a},$$

or

$$g = \frac{1}{2}\sqrt{2\pi}.$$

We can give an equation a label so that we can refer to it later.

$$E = -J \sum_{i=1}^N s_i s_{i+1}, \quad (2)$$

Equation (2) expresses the energy of a configuration of spins.[2] If we use the amsmath package, we could write Eq. (2).

We can define our own macros to save typing. For example, suppose that we introduce the macros:

```
\newcommand{\lb}{\langle}
\newcommand{\rb}{\rangle}
```

Then we can write the average value of  $x$  as

```
\begin{equation}
\lb x \rb = 3
\end{equation}
```

The result is

$$\langle x \rangle = 3. \quad (3)$$

Examples of more complicated equations:

$$I = \int_{-\infty}^{\infty} f(x) dx. \quad (4)$$

We can do some fine tuning by adding small amounts of horizontal spacing:

```
\, small space      \! negative space
```

as is done in Eq. (4).

We also can align several equations:

$$a = b \quad (5)$$

$$c = d, \quad (6)$$

or number them as subequations:

$$a = b \quad (7a)$$

$$c = d. \quad (7b)$$

If you have multiple lines for an equation but only want to number the whole equation, you suppress the equation numbers in any line by adding before . A slightly better way to align equations is to use the amsmath package:

$$\begin{aligned} a &= b \\ c &= d. \end{aligned}$$

Can you notice the difference? Some other examples:

$$m(T) = \begin{cases} 0 & T > T_c \\ (1 - [\sinh 2\beta J]^{-4})^{1/8} & T < T_c \end{cases} \quad (8)$$

$$\begin{aligned} \mathbf{T} &= \begin{pmatrix} T_{++} & T_{+-} \\ T_{-+} & T_{--} \end{pmatrix} \\ &= \begin{pmatrix} e^{\beta(J+B)} & e^{-\beta J} \\ e^{-\beta J} & e^{\beta(J-B)} \end{pmatrix} \end{aligned}$$

$$\sum_i \vec{A} \cdot \vec{B} = -P \int \mathbf{r} \cdot \hat{\mathbf{n}} dA = P \int \vec{\nabla} \cdot \mathbf{r} dV. \quad (7)$$

### 3 Lists

Some example of formatted lists include the following:

1. bread
  2. cheese
- Tom
  - Dick

An example of a table is given in Table 1 at the end of the manuscript and examples of how to include figures are shown in Figs. 1 and ??.

## 4 Special Symbols

### 4.1 Common Greek letters

These commands may be used only in math mode. Only the most common letters are included.

$$\alpha, \beta, \gamma, \Gamma, \delta, \Delta, \epsilon, \zeta, \eta, \theta, \Theta, \kappa, \lambda, \Lambda, \mu, \nu, \xi, \Xi, \pi, \Pi, \rho, \sigma, \tau, \phi, \Phi, \chi, \psi, \Psi, \omega, \Omega$$

## 4.2 Special symbols

The derivative is defined as

$$\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} \quad (7)$$

$$f(x) \rightarrow y \quad \text{as} \quad x \rightarrow x_0 \quad (7)$$

$$f(x) \xrightarrow{x \rightarrow x_0} y \quad (7)$$

Order of magnitude:

$$\log_{10} f \simeq n \quad (7)$$

$$f(x) \sim 10^n \quad (7)$$

Approximate equality:

$$f(x) \simeq g(x) \quad (7)$$

$\TeX$  is simple if we keep everything in proportion:

$$f(x) \propto x^3. \quad (7)$$

We can skip some space by using a command such as

```
\bigskip \medskip \smallskip \vspace{1in}
```

The space can be negative.

And it sometimes is convenient to write  $\tilde{x}$ ,  $\widetilde{xy}$ ,  $\overline{A}$ ,  $\%$ , accents: Schrödinger and  $\acute{e}$  (or any other letter  $\acute{z}$ ).

## 5 Literal text

It is desirable to print program code exactly as it is typed in a monospaced font. Use `\begin{verbatim}` and `\end{verbatim}` as in the following example:

```
public void computeArea()
{
    this.area = this.length*this.length;
    System.out.println("Area = " + this.area);
}
```

The command `\verbatiminput{programs/Square.java}` allows the file `Square.java` in the direction `programs` to be listed without changes.

## A Some dos and don'ts

1. Note the AJP style for books[1] and articles.[3]
2. Also note the American convention of the positions of citations.
3. Do not skip a line before `\begin{equation}` or after `\end{equation}`.

## References

- [1] Helmut Kopka and Patrick W. Daly, *A Guide to L<sup>A</sup>T<sub>E</sub>X: Document Preparation for Beginners and Advanced Users* (Addison-Wesley, Reading, MA, 1999), 3rd. ed.
- [2] It is necessary to process a file twice to get the counters correct. AJP does not use footnotes.
- [3] B. C. Freasier, C. E. Woodward, and R. J. Bearman, "Heat capacity extrema on isotherms in one-dimension: Two particles interacting with the truncated Lennard-Jones potential in the canonical ensemble," *J. Chem. Phys.* **105**, 3686–3690 (1996).

## Tables

Tables are a little difficult until you get the knack.  $\LaTeX$  automatically calculates the width of the columns. Tables should be placed at the end of the manuscript.

lattice	$d$	$q$	$T_{\text{mf}}/T_c$
square	2	4	1.763
triangular	2	6	1.648
diamond	3	4	1.479
simple cubic	3	6	1.330
bcc	3	8	1.260
fcc	3	12	1.225

Table 1: Comparison of the mean-field predictions for the critical temperature of the Ising model with exact results and the best known estimates for different spatial dimensions  $d$  and lattice symmetries.

## Figures

It is easy to include jpg or pdf or png files (see Figure 1). We can make figures bigger or smaller by scaling them. Figure 2 has been scaled by 40% to make it smaller. Figures should be placed at the end of the manuscript and sent as separate files. It also is possible to include pdf files.




Figure 1: Show me a sine.




Figure 2: Plot of the Lennard-Jones potential  $u(r)$ . The potential is characterized by a length  $\sigma$  and an energy  $\epsilon$ . This potential is applied in Ref. [3].